

CHILDHOOD IMMUNISATION SCHEDULE

IMMUNISATION
PROTECTS YOUR CHILD FROM SERIOUS ILLNESSES

Age of Child	Vaccines Needed	How and Where it is given
At Birth	BCG: Bacillus Calmette Guerin	Right Arm
	OPV (0): Oral Polio Vaccine	Drops By Mouth
6 Weeks	OPV (1): Oral Polio Vaccine	Drops By Mouth
	RV (1): Rotavirus Vaccine	Liquid By Mouth
	DTaP- IPV - HiB- HBV (1): Diphtheria, Tetanus, Accellular Pertussis, Inactivated Polio vaccine, Haemophilus influenzae type b and Hepatitis B Combined	Injection: Left Thigh
	PCV (1): Pneumococcal Conjugated Vaccine	Injection: Right Thigh
10 Weeks	DTaP- IPV - HiB- HBV (2): Diphtheria, Tetanus, Accellular Pertussis, Inactivated Polio vaccine, Haemophilus Influenzae Type B and Hepatitis B Combined	Injection: Left Thigh
14 Weeks	RV (2): Rotavirus Vaccine	Liquid By Mouth
	DTaP- IPV - HiB- HBV (3): Diphtheria, Tetanus, Accellular Pertussis, Inactivated Polio vaccine, Haemophilus Influenzae Type B and Hepatitis B Combined	Injection: Left Thigh
	PCV (2): Pneumococcal Conjugated Vaccine	Injection: Right Thigh
6 Months	Measles Vaccine (1)	Injection: Left Thigh
9 Months	PCV (3): Pneumococcal Conjugated Vaccine	Injection: Right Thigh
12 Months	Measles Vaccine (1)	Injection: Right Arm
18 Months	DTaP- IPV - HiB- HBV (4): Diphtheria, Tetanus, Accellular Pertussis, Inactivated Polio vaccine, Haemophilus Influenzae Type B and Hepatitis B Combined	Injection: Left Arm
6 Years	Td Vaccine: Tetanus and reduced strength of Diphtheria Vaccine	Injection: Left Arm
12 Years	Td Vaccine: Tetanus and reduced strength of Diphtheria Vaccine	Injection: Left Arm



CITY OF CAPE TOWN
ISIXEKO SASEKAPA
STAD KAAPSTAD

Making progress possible. Together.